


C O S M O C O R E

Core-Shell Particle for UHPLC resolution

2.6C₁₈

2.6CHOLESTER

2.6PBR




BGB GC|LC
MS|CE



COSMOCORE SERIES

◆ About Core-Shell Particles


Core-shell particles consist of a nonporous core inside a porous shell. By using these core-shell particles, one can achieve sharper peaks compared to fully porous silica gel particles of the same diameter.




Schematic diagram of a silica particle

◆ Uniform Particle Size Distribution Compared to 1.7 μm Particles

Compared to fully porous particles, core-shell particles have a more uniform particle diameter, therefore core-shell particles can be packed in the column more uniformly to minimize sample diffusion.




COSMOCORE 2.6C₁₈ (200x)




Fully porous 1.7 μm particles (200x)

◆ Mass Transfer Equivalent to Fully Porous sub-2 μm Particles

Mass transfer refers to the time it takes for a sample molecule to enter and leave a particle. In general, lower mass transfer time corresponds to less diffusion and sharper peaks. Even though COSMOCORE 2.6C₁₈ has a larger particle diameter than fully porous sub-2 μm particles, the mass transfer characteristics are similar.





COSMOCORE 2.6C₁₈



Fully porous sub-2 μm particle

◆ Reduced Back Pressure and Faster Analyses

COSMOCORE 2.6C₁₈ delivers performance equivalent to sub-2 μm particles at faster flow rate and analysis time while maintaining a lower back pressure. COSMOCORE can also be used in longer column size to gain additional resolution.




Column size: 2.1 mm I.D. - 50 mm, Mobile phase: Acetonitrile / Water = 70 / 30, Temperature: 40°C, Sample: Amylbenzene



◆ Same performance and lower back pressure compared to sub-2 µm particles

• Reduced Back Pressure

COSMOCORE 2.6C₁₈ maintains the same performance as sub-2 µm particles with half the back pressure.



◆ Specifications

	2.6C ₁₈	2.6Cholester	2.6PBr
Silica gel		Core-shell type silica	
Average particle size		2.6 µm	
Core average diameter		1.6 µm	
Average pore size		approx. 90 Å	
Specific surface area		approx. 150 m ² /g	
Stationary phase	 Octadecyl group	 Cholesteryl Group	 Pentabromobenzyl Group
Main interaction	Hydrophobic interaction	Hydrophobic interaction Molecular shape selectivity	Hydrophobic interaction Dispersion force
Endcapping treatment		Yes	
Usable pH range	1.5-10		2-7.5
Maximum pressure		60MPa	
Page	P4	P8	P11



COSMOCORE 2.6C₁₈

◆ Features

- Ultra-high performance LC results with conventional HPLC equipment
- Same number of theoretical plates as sub-2 µm columns with half the back pressure
- Increased loading capacity
- Excellent pH stability (1.5-10)

◆ Excellent pH Stability

Under accelerated pH 10.4, 40°C stability test, COSMOCORE C₁₈ shows superior stability compared with other core-shell C₁₈ phases.




Column size: 2.1 mm x 100 mm, Mobile phase: 0.35% Ammonium hydroxide/ Acetonitrile = 90/10 (pH 10.4), Sample: Caffeine 0.05 mg/ml, Injection volume: 1µl
Flow rate: 0.4 ml/min., Temperature: 40 °C

◆ Higher Retention and Loading Capacity than Competitors' Core-Shell Columns


• Retention

COSMOCORE 2.6C₁₈ features high retention and better separation compared to other core-shell columns.



• Higher Loading Capacity

In general, core-shell particles have less surface area and less bonded phase than fully porous particles. COSMOCORE 2.6C₁₈, however, with a large amount of bonded phase, exhibits both high retention and high loading capacity. When injecting 5-10 times more sample volumes in trace component analysis, COSMOCORE minimizes peak splitting due to sample overloading.




◆ Sharp Peaks with Many Types of Compounds


COSMOCORE 2.6C₁₈ features a special end capping treatment that effectively shields residual silanol groups, yielding sharp peaks for basic compounds and metal coordination complexes.

Basic Compounds


• Amitriptyline




• Haloperidol



• Procainamide




• Caffeine




◆ Sharp Peaks with Many Types of Compounds (Continued)

- Ciprofloxacin




Metal Coordination Complexes

- Oxine-Copper




◆ Low Bleed - Suitable for LC-MS

COSMOCORE 2.6C₁₈ has low column bleed and consequently low MS noise level.




Differences between Lots

COSMOCORE 2.6C₁₈ has great lot-to-lot reproducibility.



Durability Test

COSMOCORE 2.6C₁₈ exhibits high durability. Even after analyzing amitriptyline 1000 times, there is no significant deterioration in retention or peak shape.





COSMOCORE 2.6CHOLESTER

- Cholesterol-bonded reversed-phase core-shell column
- Usable under the same conditions as C₁₈ columns
- Better selectivity for cis-trans isomers, polyphenols, and natural products

◆ Comparison with C₁₈


COSMOCORE 2.6Cholester offers improved separation for cis-trans isomers than C18 under typical reversed-phase mobile phase.

COSMOCORE Application Data

Column: 2.1mmI.D.-100mm
Column size: 2.1mmI.D.-100mm
Mobile phase: Methanol
Flow rate: 0.4 ml/min
Temperature: 40°C
Detection: UV280nm

Sample: 1; Vitamin K₂ (0.50mg/ml)
2; trans-Vitamin K₁ (0.50mg/ml)
3; cis-Vitamin K₁

Inj. Vol: 0.5μl





Competitor A
(Fully porous, 1.7μm, C₁₈)

Competitor B
(Core-Shell, 2.6μm, C₁₈)

COSMOCORE
2.6C₁₈

COSMOCORE
2.6Cholester



NACALAI TESQUE, INC

◆ Separation Properties

COSMOCORE 2.6Cholester has about the same hydrophobicity as C₁₈. It is not necessary to change the analytical conditions when replacing C₁₈ Columns with COSMOCORE 2.6Cholester.

COSMOCORE Application Data

Column: 2.1mmI.D.-100mm
Column size: 2.1mmI.D.-100mm
Mobile phase: Acetonitrile/ H₂O = 50/50
Flow rate: 0.4 ml/min
Temperature: 40°C
Detection: UV254nm

Sample: 1; Uracil
2; Methyl Benzoate
3; Toluene
4; Naphthalene


Inj. Vol: 0.25μl

Competitor A
(Fully porous, 1.7μm, C₁₈)

Competitor B
(Core-Shell, 2.6μm, C₁₈)

COSMOCORE
2.6C₁₈


COSMOCORE
2.6Cholester



NACALAI TESQUE, INC

Molecular Shape Selectivity


COSMOCORE 2.6Cholester has excellent shape selectivity due to its structural rigidity. COSMOCORE 2.6Cholester retains planar triphenylene longer than non planar o-terphenyl.




Applications

- Separation of Natural Compounds (Tomato components)


Lycopene exists in many cis-trans isomers. COSMOCORE 2.6Cholester separates these compounds better than C₁₈.



- Separation of Catechin




- Apply to LC/MS




◆ Scaling Up from Analytical to Preparative Separation

COSMOCORE 2.6Cholester has the same functional group as the fully-porous COSMOSIL Cholester 5 µm silica packing material. Therefore, the separation pattern in COSMOCORE 2.6Cholester is maintained when scaling up to preparative separation with 5 µm COSMOSIL Cholester.

COSMOCORE Application Data



◆ References to COSMOSIL Cholesterol-Bonded Stationary Phase



◆ Cholester Series Application Notebook and Reference List

Cholester is a silica-based reversed-phase column using cholesterol as the bonded phase. It has similar hydrophobicity to C₁₈ (ODS) and, with superior stereoselectivity, it is suitable for compounds with similar hydrophobicity but slightly different molecular shape. Designed for ease of use, Cholester can be used under solvent and other analytical conditions identical to ODS. This application notebook includes over 80 applications and 88 literature references, and it features our fully porous and core-shell packing materials.

COSMOSIL
COSMOSIL/COSMOCORE Cholester Series Application Notebook and Reference List

Cholester Reference List

The number of COSMOSIL Cholester references
The total number of COSMOSIL Cholester references is 100. Individual year: 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014.

Cholester Applications

Cholester Applications (2014)

Cholester Applications (2013)

Cholester Applications (2012)

Cholester Applications (2011)

Cholester Applications (2010)

Cholester Applications (2009)

Cholester Applications (2008)

Cholester Applications (2007)

Cholester Applications (2006)

Cholester Applications (2005)



COSMOCORE 2.6PBr

- Separate hydrophilic compounds under reversed-phase conditions
- Retain hydrophilic compounds longer than C₁₈
- Greater sample loading capacity than HILIC
- High performance similar to sub-2 µm particles with lower back pressure


Suitable Samples

- Hydrophilic compounds
- Nucleic acids and derivatives
- Surfactants
- Glycosides
- Peptides


Separation of Hydrophilic Compounds (low retention on C₁₈)

COSMOSIL PBr retains hydrophilic compounds stronger than C₁₈ columns under the same reversed-phase conditions.

Nucleic Acid Metabolites




Malonyl CoA, CoA, Acetyl CoA



◆ Separation of Hydrophilic Compounds (low retention on C₁₈) (continued)


• Water-Soluble Vitamins




◆ Separation of Hydrophilic Compounds (compounds with similar hydrophobicity)

COSMOCORE 2.6PBr can separate compounds with similar hydrophobicity, utilizing several kinds of molecular interactions, including dispersion force generated by the bromine atoms.


• ATP, ADP, AMP



• Vitamin B3 (Nicotinic Acid, Nicotinamide)




• Thiamine Pyrophosphate, Thiamine




◆ Separation of Hydrophilic Compounds (Glycosides)

Glycosides with identical aglycones but different glycosyl groups can also be separated.

• UDP Glycosides



• Arbutin and Hydroquinone



◆ Separation Mechanism

► Dispersion force (instantaneous dipole-induced dipole force)

London dispersion force is a weak intermolecular force that results from dipoles temporarily induced by random unsymmetrical electron positions in two adjacent atoms, also known as instantaneous dipole-induced dipole force. It is present in all molecules regardless of polarity. Compounds with high polarizability have stronger dispersion force.

► Compounds with stronger dispersion force

- Larger and heavier molecules
- Molecules with larger and heavier atoms (e.g. from weakest to strongest in halogens, F₂, Cl₂, Br₂, and I₂)
- Molecules with delocalized electrons and resonance (e.g. aromatic compounds)




COSMOSIL PBr column is packed with pentabromobenzyl-bonded silica that enables separation by dispersion force interaction.

◆ Difference between Methanol and Acetonitrile Mobile Phase

Comparison of Separation Ability


Acetonitrile is frequently used in HPLC to reduce backpressure. However, the π electrons in acetonitrile interfere with the dispersion force interaction between the sample and the stationary phase. Therefore, we recommend using methanol as the organic solvent.



Difference of Methanol and Acetonitrile Mobile Phase (continued)

Comparison of Pressure

In general, methanol/water mobile phases exhibit significantly higher pressure than acetonitrile/water. However, the pressure never exceeded 30 MPa in our test of a 2.1 mm x 100 mm column. Since COSMOSORE 2.6PBr can withstand 60 MPa, methanol-based mobile phases are usable even with UHPLC.



Column : COSMOCORE 2.6PBr (2.1mmI.D.-100mm)
 Mobile Phase : See graph
 Temperature : 40°C
 Flow rate : 0.4 ml/min

Comparison to HILIC

Both columns are suitable for analysis of hydrophilic compounds, but they have different properties.

	PBr	HILIC
Separation Mode	Reversed Phase	Hydrophilic Interaction (HILIC)
Features	<ul style="list-style-type: none"> Simple mobile phase conditions compared to HILIC. Strong eluent: organic (methanol) Weak eluent: water Low peak distortion with water-based samples; usable with large injection volumes of dilute samples. Some hydrophilic compounds are not retained well. 	<ul style="list-style-type: none"> Retains hydrophilic compounds that would not be retained by C₁₈ columns. Strong eluent: water Weak eluent: organic (acetonitrile) Peak distortion occurs with large volumes of samples dissolved in water.




INSTRUMENT SETTINGS AND COMPATIBILITY

When Using with a Conventional (non-UHPLC) Instrument

COSMOCORE 2.6C₁₈ is designed for use with UHPLC instruments. In addition, due to its low backpressure, it can be used with conventional instruments. However, it is necessary to change the following settings.


1. Detector Response Time

Because UHPLC analyses are done at high flow rates, a slow response time can adversely affect peak shape. We recommend setting the response time to 0.1 sec or less.



2. Other Instrument Parameters

UHPLC is more vulnerable to the effects of dead volume than conventional chromatography. When using a 2.1 mm I.D. column, please use a semi-micro detector cell, injector, and piping (0.1 mm).



• Fittings and Adapters

COSMOCORE columns use the same connectors as Waters UPLC® (UHPLC) columns. This is different from our conventional COSMOSIL columns, which use the conventional Waters HPLC-compatible connectors. (UPLC® is a registered trademark of Waters Corporation.)

1. Differences between End Fitting

Connection type		Column	
	HPLC (COSMOSIL)	UHPLC (COSMOCORE)	
Instrument	HPLC	No adapter required	Adapter required
	UHPLC	Adapter required	No adapter required

HPLC: Conventional Waters-compatible connector

UHPLC: Waters UPLC®-compatible connector


Connector shape	HPLC	UHPLC

The length of tubing that extends from the ferrule differs from HPLC to UHPLC.

2. COSMOCORE-Compatible Fittings

1) UHPLC Instrument Fittings

No adapter needed; just connect as-is.




2) HPLC Instrument Fittings

An adapter or movable (high-pressure) fitting is required to connect the fittings to the column. See the examples for different fittings below.


• SUS Ferrules (HPLC) Fixed on the Tubing

The column can be connected by using an adapter (SUS union + tubing fixed to UHPLC length).



• PEEK Fittings

PEEK fittings do not fix the length of tubing at the end, so they can be used with both types of column. However, please be cautious of their pressure tolerance.





ORDERING INFORMATION

COSMOCORE Series

Column Size (mm I.D. x mm)	2.6C ₁₈ Product No.	2.6Cholester Product No.	2.6PBr Product No.
2.1 x 30	12632-31	12858-91	13692-21
2.1 x 50	12631-41	12859-81	13693-11
2.1 x 75	12630-51	12860-41	13694-01
2.1 x 100	12614-71	12861-31	13695-91
2.1 x 150	12612-91	12862-21	13697-71
3.0 x 30	12611-01	12863-11	13698-61
3.0 x 50	12609-51	12864-01	13699-51
3.0 x 75	12608-61	12866-81	13700-01
3.0 x 100	12607-71	12867-71	13701-91
3.0 x 150	12602-21	12868-61	13703-71

Column Size (mm I.D. x mm)	2.6C ₁₈ Product No.	2.6Cholester Product No.	2.6PBr Product No.
4.6 x 30	12601-31	12869-51	13705-51
4.6 x 50	12600-41	12870-11	13712-51
4.6 x 75	12599-91	12871-01	13714-31
4.6 x 100	12598-01	12872-91	13715-21
4.6 x 150	12597-11	12873-81	13719-81
4.6 x 250	12596-21	12875-61	13734-71

- COSMOCORE's connector is the same type as Waters UPLC® Columns.
- Other sizes may be available.
- These phases are also available as fully porous particles. For more information, please see <http://www.nacalai.co.jp/URL/?P=TOP>

Adapter List

Name	Description	Product Number	PKG Size
Low & Zero Dead Volume Union	Material: SUS Bore diameter: 0.35 mm	P0402	1 PKG
COSMOSIL Column Connecting Tube (0.1 mm I.D.)	I.D.: 0.1 mm	12570-41	1 PKG
COSMOSIL Column Connecting Tube (0.25 mm I.D.)	I.D.: 0.25 mm	37843-69	1 PKG

UHPLC-Compatible Prefilter

Product Name	In	Out	Contents	Product Number	PKG Size
U-Fil UHPLC-compatible prefILTER	UHPLC	UHPLC	Filter: 0.5 µm	12571-31	1 PKG
	HPLC	UHPLC	Tubing connecting diameter: 1/16	12572-21	1 PKG
U-Fil replacement filter	-	-	Filter: 0.5 µm, Material: SUS316L	28150	5 units / Pack

For research use only, not intended for diagnostic or drug use.