

Packing Materials for Column Chromatography


Open column chromatography is an excellent and easy technique for large-scale preparation and purification at low cost. COSMOSIL offers both normal and reversed phase packing materials based on totally porous spherical silica, which provides higher separation, less pressure and higher reproducibility than irregular silica.

Material Characteristics

Packing Material	C ₁₈ -OPN	C ₁₈ -PREP	SL-II-PREP	Silica Gel 60 (neutral)	
Silica Gel	High Purity Porous Spherical Silica				
Average Particle Size	75, 140 µm	40, 75, 140 µm	75, 140 μm		
Average Pore Size	approx. 120 Å		approx. 60 Å		
Specific Surface Area	approx. 300 m ² /g		approx. 300 m ² /g approx. 500 m ² /g		
Stationary phase	Octadeo	Octadecyl Group		None	
Carbon content	-	approx. 19%	approx. 0%		
End-capping treatment	None	Treated	-		
Llooful rongo	Open column chromatography / Flash column chromatography				
Useful range	Reversed phase	e chromatography	Normal phase chromatography		




Selection Guide



COSMOSIL C₁₈-OPN

- \cdot A new "Water-Wet" C₁₈ packing material for reversed phase open column chromatography
- \cdot Enables large-scale preparation and purification at low cost

Characteristic



The external surface of the C18-OPN gel is coated with hydrophilic groups.



Packing materials in water Left : C18 OPN provides good resolution Right : Conventional product float up

Conventional reversed phase C_{18} packing materials are restricted to about 30-50% water in the mobile phase. The COSMOSIL C_{18} -OPN is a new "Water-Wet" C_{18} packing material developed for reversed phase open column chromatography. The C_{18} -OPN material can be used in 100% aqueous eluents.

Application Data


· Separation of hydrophilic compounds in aqueous solution

COSMOSIL Application Data 75C₁₈-OPN Column: 20mmI.D.-250mm Column size: Mobile phase: Methanol/ $H_2O=30/70$ Flow rate: 0.2 ml/min Temperature: Detection: Room temperature UV254nm Sample: 1; Theobromine (100mg) 2; Theophylline (100mg) 30 60 90 120 150 Time (min) NACALAI TESQUE, INC

In reversed phase chromatography, hydrophilic compounds such as Theobromine and Theophylline could be separated under low concentration of organic solvent. The figure shows they are clearly separated by reversed open column chromatography with 70% of water.

• Separation of *p*-Cresol and *p*-Ethylphenol by normal and reversed phase mode

Since the structural difference between *p*-Cresol and *p*-Ethylphenol is only one methyene group. It is difficult to separate such samples under normal phase condition. On the other hand, the samples are clearly separated under reversed phase condition with COSMOSIL C_{18} -OPN packing material.




COSMOSIL C₁₈-PREP

- · Standard reversed phase packing material for open chromatography
- Enables large-scale preparation and purification at low cost


Application Data

• Vitamin E

Natural Compounds



Basic Compounds



Highly Polar Compounds

· Reproducibility and washing methods

Wash the COSMOSIL C₁₈-OPN packing material with tetrahydrofuran, chloroform or other solvents to remove the impurities. This packing material has excellent reproducibility and can be used repeatedly.

"CAUTION"

Do not wash with basic solvents of pH 7 or more which will dissolve the silica gel or pH 2 or less which will cleave the C_{18} stationary phase. Dry the packing material at 50°C or less. See end of this chapter for packing method.

COSMOSIL SL-II-PREP


· Standard packing materials for normal phase chromatography

· Ultra pure silica gel packing material more than 99.99% purity


Comparison

Highly purified silica gel of COSMOSIL SL-II-PREP enables separation of metal coordination compounds without adsorption.

Metal Coordination Compounds



Organic Acid and Amide



Silica Gel 60 (spherical · neutral)


• The pH of Silica Gel is adjusted to neutral

Suitable for the separation of pH sensitive compounds

Comparison with conventional silica gel

Since conventional silica gels are weakly acidic, some pH sensitive compounds may be decomposed during the purification by column chromatography with the acidic silica gels. The pH of Silica Gel 60 (spherical • neutral) is adjusted to nearly neutral for the separation of not only pH sensitive compounds but also new compounds that the physical properties are still unknown.

Purification of Acetal



Ordering Information

· COSMOSIL C₁₈-OPN

· COSMOSIL C₁₈-PREP

10			
Product Name	Particle Size	PKG Size	Product Number
COSMOSIL 75C ₁₈ -OPN		100 g	37842-66
	75 µm	500 g	37842-95
		1 kg	37842-11
COSMOSIL 140C ₁₈ -OPN		100 g	37878-16
	140 µm 🤤	500 g	37878-45
		1 kg	37878-61

COSMOSIL SL-II PREP

Product Name	Particle Size	PKG Size	Product Number
COSMOSIL 75SL-II-PREP		100 g	38012-64
	75 µm	500 g	38012-35
		1 kg	38012-51
COSMOSIL 140SL-II-PREP	140 µm	100 g	38013-54
		1 kg	38013-41

· Silica Gel (spherical · neutral)

	,		
Product Name	Particle Size	PKG Size	Product Number
Silica Gel 60	75 µm	500 g	30511-35
(spherical • neutral)		1 kg	30511-51
for column chromatograph	140 µm	500 g	30518-65
		1 kg	30518-81

Product Name	Particle Size	PKG Size	Product Numb
COSMOSIL 40C ₁₈ -PREP		100 g	37932-86
	40 µm	500 g	37932-15
		1 kg	37932-31
COSMOSIL 75C18-PREP		100 g	37933-76
	75 µm	500 g	37933-05
		1 kg	12061-11
COSMOSIL 140C ₁₈ -PREP		100 g	37934-66
	140 µm	500 g	37934-95

For research use only, not intended for diagnostic or drug use.



NACALAI TESQUE, INC.

umber -86

37934-11

1 kg

Nijo Karasuma, Nakagyo-ku, Kyoto 604-0855 JAPAN TEL

- : +81-(0)75-251-1730 : +81-(0)75-251-1763
- FAX Website : www.nacalai.com

: info.intl@nacalai.com E-mail